

---

**LOCAL CONVERGENCE OF A MULTI-POINT JARRATT-TYPE METHOD IN  
BANACH SPACE UNDER WEAK CONDITIONS**

IOANNIS K. ARGYROS\* AND SANTHOSH GEORGE

Ioannis K. Argyros, Cameron University, Department of Mathematical Sciences, Lawton, OK 73505,  
USA, Email: iargyros@cameron.edu

Department of Mathematical and Computational Sciences, National Institute of Technology Karnataka,  
India-757 025, Email: sgeorge@nitk.ac.in

---

**ABSTRACT.** We present a local convergence analysis of a multi-point Jarratt-type method of high convergence order in order to approximate a solution of a nonlinear equation in a Banach space. Our sufficient convergence conditions involve only hypotheses on the first Fréchet-derivative of the operator involved. In contrast to earlier studies using hypotheses up to the third Fréchet-derivative [26]. Numerical examples are also provided in this study.

**KEYWORDS :** Jarratt-type methods, Banach space, Local Convergence, Fréchet-derivative.

**AMS Subject Classification:**65H10, 65D99.

1. INTRODUCTION

In this study we are concerned with the problem of approximating a solution  $x^*$  of the nonlinear equation

$$F(x) = 0, \tag{1.1}$$

where  $F$  is a Fréchet-differentiable operator defined on a subset  $D$  of a Banach space  $X$  with values in a Banach space  $Y$ .

Many problems in computational sciences and other disciplines can be brought in a form like (1.1) using mathematical modelling [3]. The solutions of equation (1.1) can rarely be found in closed form. That is why solutions of equation (1.1) are approximated by iterative methods. In particular, the practice of Numerical Functional Analysis for finding such solution is essentially connected to Newton-like methods [1]-[27]. The study about convergence matter of iterative procedures is usually based on two types: semi-local and local convergence analysis. The semi-local convergence matter is, based on the information around an initial point, to give conditions ensuring the convergence of the iterative procedure; while the local one is, based on the information around a solution, to find estimates of the

---

\* Corresponding author.  
Email address : iargyros@cameron.edu.  
Article history : Received April 8, 2014. Accepted February 18, 2016.

radii of convergence balls. There exist many studies which deal with the local and semilocal convergence analysis of Newton-like methods such as [1]-[27].

We present a local convergence analysis for the multi-point Jarratt-type method defined for each  $n = 0, 1, 2, \dots$  by

$$\begin{aligned} u_n &= x_n - F'(x_n)^{-1}F(x_n), \\ y_n &= u_n + \frac{1}{3}F'(x_n)^{-1}F(x_n), \\ z_n &= u_n + B_n F(x_n), \\ x_{n+1} &= z_n - A_n F(z_n), \end{aligned} \tag{1.2}$$

where  $x_0 \in D$  is an initial point,  $J_n = (6F'(y_n) - 2F'(x_n))^{-1}(3F'(y_n) + F'(x_n))$ ,  $B_n = (I - J(x_n))F'(x_n)^{-1}$ , and  $A_n = \frac{3}{2}J_n F'(y_n)^{-1} + (I - \frac{3}{2}J_n)F'(x_n)^{-1}$ .

A semilocal convergence analysis was given in [26] but the operator  $A_n$  was defined by

$$\overline{A}_n = \frac{3}{2}F'(y_n)^{-1}J_n + F'(x_n)^{-1}(I - \frac{3}{2}J_n).$$

That is, their method is defined by

$$\begin{aligned} u_n &= x_n - F'(x_n)^{-1}F(x_n), \\ y_n &= u_n + \frac{1}{3}F'(x_n)^{-1}F(x_n), \\ z_n &= u_n + B_n F(x_n), \\ x_{n+1} &= z_n - \overline{A}_n F(z_n). \end{aligned} \tag{1.3}$$

Notice that for two linear operators  $Q_1$  and  $Q_2$  we have that  $Q_1Q_2 \neq Q_2Q_1$ , so  $A_n \neq \overline{A}_n$  in general. The fifth order of convergence of method (1.3) was established in [26]. These results were given in a non-affine invariant form. However, the results obtained in our paper are given in affine invariant form. The sufficient semilocal convergence conditions (given in affine invariant form) used in [26] are (C):

(C<sub>1</sub>): There exists  $F'(x_0)^{-1} \in L(Y, X)$  and  $\|F'(x_0)^{-1}\| \leq \beta$ ;

(C<sub>2</sub>):

$$\|F'(x_0)^{-1}F(x_0)\| \leq \beta_1;$$

(C<sub>3</sub>):

$$\|F'(x_0)^{-1}F''(x)\| \leq \beta_2 \quad \text{for each } x \in D;$$

(C<sub>4</sub>):

$$\|F'(x_0)^{-1}F'''(x)\| \leq \beta_3 \quad \text{for each } x \in D;$$

(C<sub>5</sub>):

$$\|F'(x_0)^{-1}(F'''(x) - F'''(y))\| \leq w(\|x - y\|) \quad \text{for each } x, y \in D$$

where  $w(s)$  is a nondecreasing continuous real function for  $s > 0$  with  $w(0) \geq 0$ .

(C<sub>6</sub>): there exists a non-negative real function  $\phi \in C[0, 1]$ , with  $\phi(t) \leq 1$ , such that  $w(ts) \leq \phi(t)w(s)$  for each  $t \in [0, 1]$ ,  $s \in (0, \infty)$ .

Similar conditions have been used by other authors [1]-[27], on other high convergence order methods. The corresponding conditions for the local convergence analysis are given by simply replacing  $x_0$  by  $x^*$  in the preceding (C) conditions.

These conditions however are very restrictive. As an academic example, let us define function  $F$  on  $X = [-\frac{5}{2}, \frac{1}{2}]$  by

$$F(x) = \begin{cases} x^3 \ln x^2 + x^5 - x^4, & x \neq 0 \\ 0, & x = 0 \end{cases}.$$

Then, obviously, e.g. function  $F$  cannot satisfy condition  $(\mathcal{C}_4)$  since  $F'''(x) = 6 \ln x^2 + 60x^2 - 24x + 22$  is unbounded on  $D$ . In the present paper we only use hypotheses on the first Fréchet derivative (see conditions (2.9)-(2.12)). This way we expand the applicability of method (1.2).

The paper is organized as follows. The local convergence of method (1.2) is given in Section 2, whereas the numerical examples are given in the concluding Section 3.

## 2. LOCAL CONVERGENCE ANALYSIS

We present the local convergence analysis of method (1.2) in this section. Denote by  $U(v, \rho)$ ,  $\bar{U}(v, \rho)$  the open and closed balls, respectively, in  $X$  of center  $v$  and radius  $\rho > 0$ .

Let  $L_0 > 0$ ,  $L > 0$  and  $M > 0$  be given parameters. It is convenient for the local convergence analysis of method (1.2) that follows to define functions on the interval  $[0, \frac{1}{L_0})$  by

$$\begin{aligned} g_1(r) &= \frac{Lr}{2(1-L_0r)}, \\ g_2(r) &= g_1(r) + \frac{M}{3(1-L_0r)}, \\ g_3(r) &= \frac{3L_0(1+g_2(r))r}{2(1-L_0r)}, \\ g_4(r) &= g_1(r) + \frac{3ML_0(1+g_2(r))r}{4(1-g_3(r))(1-L_0r)^2}, \\ g_5(r) &= \left[1 + \frac{3}{4}\left(1 + \frac{1}{1-g_3(r)}\right)\frac{M}{1-L_0g_2(r)r} \right. \\ &\quad \left. + \left(1 + \frac{3}{4}\left(1 + \frac{1}{1-g_3(r)}\right)\right)\frac{M}{1-L_0r}\right]g_4(r). \end{aligned}$$

We have that

$$\begin{aligned} g_2(r) &= g_1(r) + \frac{M}{3(1-L_0r)} \\ &= \frac{Lr}{2(1-L_0r)} + \frac{M}{3(1-L_0r)}. \end{aligned}$$

Hence, if

$$r_2 = \frac{3-M}{3(\frac{L}{2} + L_0)}$$

and

$$M < 3, \tag{2.1}$$

then, we have that

$$0 < g_2(r) < 1 \text{ and } 0 < g_1(r) < 1 \text{ for each } r \in [0, r_2). \tag{2.2}$$

Notice that

$$r_2 < r_R := \frac{2}{3L} < r_A := \frac{1}{\frac{L}{2} + L_0} \text{ for } L_0 < L \tag{2.3}$$

and

$$r_2 < r_A = r_R \text{ for } L_0 = L. \quad (2.4)$$

Function  $g_3$  can be written as

$$\begin{aligned} g_3(r) &= \frac{3}{2} \frac{L_0 r}{1 - L_0 r} \left( 1 + \frac{Lr}{2(1 - L_0 r)} + \frac{M}{3(1 - L_0 r)} \right) \\ &= \frac{L_0 r}{4(1 - L_0 r)^2} (6(1 - L_0 r) + 3Lr + 2M). \end{aligned}$$

Define polynomial  $p_3$  by

$$p_3(r) = L_0 r (6(1 - L_0 r) + 3Lr + 2M) - 4(1 - L_0 r)^2.$$

We have that  $p_3(0) = -4 < 0$  and  $p_3(\frac{1}{L_0}) = \frac{3L}{L_0} + 2M > 0$ . It then follows from the intermediate value theorem that polynomial  $p_3$  has roots in the interval  $(0, \frac{1}{L_0})$ . Let us denote by  $r_3$  the smallest such root. Then, we have that

$$0 < p_3(r) < 1 \text{ and } 0 < g_3(r) < 1 \text{ for each } r \in [0, r_3]. \quad (2.5)$$

By some algebraic manipulation we see that function  $g_5$  can be written as

$$g_5(r) = \frac{N(r)}{D(r)},$$

where

$$\begin{aligned} N(r) &= [4(1 - L_0 g_2(r)r)(1 - g_3(r))(1 - L_0 r) + 3M(1 - g_3(r))(1 - L_0 r) \\ &\quad + 3M(1 - L_0 r) + 7M(1 - L_0 g_2(r)r)(1 - g_3(r)) + 3M(1 - L_0 g_2(r)r)] \\ &\quad \times [4(1 - g_3(r))(1 - L_0 r)^2 g_1(r) + 3ML_0(1 + g_2(r)r)] \end{aligned}$$

and

$$D(r) = 16(1 - g_3(r))^2 (1 - L_0 r)^3 (1 - L_0 g_2(r)r).$$

Moreover, define function  $g_6$  on the interval  $[0, \frac{1}{L_0})$  by

$$g_6(r) = N(r) - D(r) \quad (2.6)$$

we have that  $g_6(0) = -16(1 - \frac{L_0 M r}{3}) < 0$ , since  $L_0 r < 1$  and  $M < 3$  and  $g_6(r) \rightarrow \infty$  as  $r \rightarrow (\frac{1}{L_0})^-$ . It follows that function  $g_6$  has zeros in the interval  $(0, \frac{1}{L_0})$  (i.e., function  $g_5$  has zeros in the interval  $(0, \frac{1}{L_0})$ ). Denote by  $r_5$  the smallest such zero. Then, we have that

$$0 < g_5(r) < 1 \text{ and } 0 < g_4(r) < 1 \text{ for each } r \in [0, r_5]. \quad (2.7)$$

Finally, set

$$r^* = \min\{r_2, r_3, r_5\}. \quad (2.8)$$

Then, clearly (2.2), (2.5) and (2.7) hold for each  $r \in [0, r^*]$  (provided that (2.1) and (2.6) also hold).

Next, we present the local convergence analysis of method (1.2).

**THEOREM 2.1.** Let  $F : D \subseteq X \rightarrow Y$  be a Fréchet-differentiable operator. Suppose that there exist  $x^* \in D$ , parameters  $L_0 > 0, L > 0$  and  $0 < M < 3$  such that for all  $x \in D$

$$F(x^*) = 0, \quad F'(x^*)^{-1} \in L(Y, X), \quad (2.9)$$

$$\|F'(x^*)^{-1}(F(x) - F(x^*))\| \leq L_0 \|x - x^*\|, \quad (2.10)$$

$$\|F'(x^*)^{-1}(F(x) - F(x^*) - F'(x)(x - x^*))\| \leq \frac{L}{2} \|x - x^*\|^2, \quad (2.11)$$

$$\|F'(x^*)^{-1}F'(x)\| \leq M \quad (2.12)$$

and

$$\bar{U}(x^*, r^*) \subseteq D,$$

where  $r^*$  is given by (2.8). Then, sequence  $\{x_n\}$  generated by method (1.2) for  $x_0 \in U(x^*, r^*)$  is well defined, remains in  $U(x^*, r^*)$  for each  $n = 0, 1, 2, \dots$  and converges to  $x^*$ . Moreover, the following estimates hold for each  $n = 0, 1, 2, \dots$ ,

$$\|u_n - x^*\| \leq g_1(\|x_n - x^*\|)\|x_n - x^*\| < \|x_n - x^*\| < r^*, \quad (2.13)$$

$$\|y_n - x^*\| \leq g_2(\|x_n - x^*\|)\|x_n - x^*\| < \|x_n - x^*\|, \quad (2.14)$$

$$\|z_n - x^*\| \leq g_4(\|x_n - x^*\|)\|x_n - x^*\| < \|x_n - x^*\|, \quad (2.15)$$

$$\frac{3}{2}\|F'(x_n)^{-1}(F'(y_n) - F'(x_n))\| \leq g_3(\|x_n - x^*\|) < 1 \quad (2.16)$$

and

$$\|x_{n+1} - x^*\| \leq g_5(\|x_n - x^*\|)\|x_n - x^*\| < \|x_n - x^*\|, \quad (2.17)$$

where the “ $g$ ” functions are defined above Theorem 2.1.

**Proof.** Using (2.10), the definition of  $r^*$  and the hypothesis  $x_0 \in U(x^*, r^*)$  we get that

$$\|F'(x^*)^{-1}(F(x_0) - F(x^*))\| \leq L_0\|x_0 - x^*\| < L_0r^* < 1. \quad (2.18)$$

It follows from (2.18) and the Banach Lemma on invertible operators [3] that  $F'(x_0)^{-1} \in L(Y, X)$  and

$$\|F'(x_0)^{-1}F'(x^*)\| \leq \frac{1}{1 - L_0\|x_0 - x^*\|} < \frac{1}{1 - L_0r^*}. \quad (2.19)$$

Hence,  $u_0, y_0$  are well defined. Using method (1.2) for  $n = 0$ , (2.19), (2.2), (2.17), (2.11), (2.12) (for  $x = x_0$ ) and the definition of functions  $g_1$  and  $g_2$ , we obtain in turn that

$$\begin{aligned} u_0 - x^* &= x_0 - x^* - F'(x_0)^{-1}F(x_0) \\ &= F'(x_0)^{-1}[-F(x_0) + F'(x_0)(x_0 - x^*)] \\ &= -F'(x_0)^{-1}F'(x^*)F'(x^*)^{-1}[F(x_0) - F(x^*) - F'(x_0)(x_0 - x^*)] \end{aligned}$$

so,

$$\begin{aligned} \|u_0 - x^*\| &\leq \|F'(x_0)^{-1}F'(x^*)\|\|F'(x^*)^{-1}[F(x_0) - F(x^*) - F'(x_0)(x_0 - x^*)]\| \\ &\leq \frac{L\|x_0 - x^*\|^2}{2(1 - L_0\|x_0 - x^*\|)} \\ &= g_1(\|x_0 - x^*\|)\|x_0 - x^*\| < \|x_0 - x^*\| < r^*, \end{aligned}$$

which shows (2.13) for  $n = 0$ . Consequently from

$$y_0 - x^* = u_0 - x^* + \frac{1}{3}F'(x_0)^{-1}F'(x^*)F'(x^*)^{-1}F(x_0),$$

we get that

$$\begin{aligned} \|y_0 - x^*\| &\leq \|u_0 - x^*\| + \frac{1}{3}\|F'(x_0)^{-1}F'(x^*)\|\|F'(x^*)^{-1}F(x_0)\| \\ &\leq g_1(\|x_0 - x^*\|)\|x_0 - x^*\| + \frac{1}{3} \frac{\int_0^1 \|F'(x^*)^{-1}F'(x^* + t(x^* - x_0))\| dt}{1 - L_0\|x_0 - x^*\|} \|x_0 - x^*\| \\ &\leq [g_1(\|x_0 - x^*\|) + \frac{M}{3(1 - L_0\|x_0 - x^*\|)}] \|x_0 - x^*\| \\ &= g_2(\|x_0 - x^*\|)\|x_0 - x^*\| < \|x_0 - x^*\| < r^*, \end{aligned}$$

which shows (2.14) for  $n = 0$ . We need a norm estimate on  $B_0$ . Let us start with an estimate on

$$\begin{aligned}
& \frac{3}{2} \|F'(x_0)^{-1}(F'(y_0) - F'(x_0))\| \\
& \leq \frac{3}{2} \|F'(x_0)^{-1}F'(x^*)\| (\|F'(x^*)^{-1}(F'(y_0) - F'(x^*))\| \\
& \quad + \|F'(x^*)^{-1}(F'(x_0) - F'(x^*))\|) \\
& \leq \frac{3}{2} \frac{L_0(\|y_0 - x^*\| + \|x_0 - x^*\|)}{1 - L_0\|x_0 - x^*\|} \\
& \leq \frac{3}{2} \frac{L_0(1 + g_2(\|x_0 - x^*\|))\|x_0 - x^*\|}{1 - L_0\|x_0 - x^*\|} \\
& = g_3(\|x_0 - x^*\|) < g_3(r^*) < 1,
\end{aligned}$$

(by (2.5)) which shows (2.16) for  $n = 0$ ,  $(I + \frac{3}{2}F'(x_0)^{-1}(F'(y_0) - F'(x_0)))^{-1} \in L(Y, X)$  and

$$\|(I + \frac{3}{2}F'(x_0)^{-1}(F'(y_0) - F'(x_0)))^{-1}\| \leq \frac{1}{1 - g_3(\|x_0 - x^*\|)}.$$

Then, we have the estimate

$$\begin{aligned}
& \frac{1}{2}(I + \frac{3}{2}F'(x_0)^{-1}(F'(y_0) - F'(x_0)))^{-1}F'(x_0)^{-1} \\
& = [2F'(x_0)(I + \frac{3}{2}F'(x_0)^{-1}(F'(y_0) - F'(x_0)))^{-1} \\
& = [3(F'(y_0) - F'(x_0)) + 2F'(x_0)]^{-1} \\
& = (3F'(y_0) - F'(x_0))^{-1}.
\end{aligned}$$

Then,  $B_0$  can now be written as

$$B_0 = \frac{3}{4}(I + \frac{3}{2}F'(x_0)^{-1}(F'(y_0) - F'(x_0)))^{-1}F'(x_0)^{-1}(F'(y_0) - F'(x_0))F'(x_0)^{-1},$$

so,

$$\begin{aligned}
\|B_0\| & \leq \frac{3}{4} \frac{1}{1 - g_3(\|x_0 - x^*\|)} \\
& \quad \times \frac{L_0(\|x_0 - x^*\| + \|y_0 - x^*\|)}{1 - L_0\|x_0 - x^*\|} \frac{M\|x_0 - x^*\|}{1 - L_0\|x_0 - x^*\|} \\
& \leq \frac{3}{4} \frac{ML_0(1 + g_2(\|x_0 - x^*\|))\|x_0 - x^*\|^2}{(1 - g_3(\|x_0 - x^*\|))(1 - L_0\|x_0 - x^*\|)^2}. \tag{2.20}
\end{aligned}$$

Moreover from (2.20) and (2.7) we get that

$$\begin{aligned}
\|z_0 - x^*\| & \leq \|u_0 - x^*\| + \|B_0F(x_0)\| \\
& \leq [g_1(\|x_0 - x^*\|) + \frac{3}{4} \frac{ML_0(1 + g_2(\|x_0 - x^*\|))\|x_0 - x^*\|}{(1 - g_3(\|x_0 - x^*\|))(1 - L_0\|x_0 - x^*\|)^2}] \|x_0 - x^*\| \\
& = g_4(\|x_0 - x^*\|)\|x_0 - x^*\| < \|x_0 - x^*\| < r^*,
\end{aligned}$$

which shows (2.15) for  $n = 0$ .

Next, we need an estimate on  $J_0$  and  $A_0$ . We have that

$$\begin{aligned}
J_0 & = \frac{1}{2}(3F'(y_0) - F'(x_0))^{-1}(3F'(y_0) - F'(x_0) + 2F'(x_0)) \\
& = \frac{1}{2}(I + 2(3F'(y_0) - F'(x_0))^{-1}F'(x_0))
\end{aligned}$$

so,

$$\|J_0\| \leq \frac{1}{2} \left(1 + \frac{1}{1 - g_3(\|x_0 - x^*\|)}\right)$$

and

$$\begin{aligned} \|A_0 F(z_0)\| &\leq \left[ \frac{3}{2} \|J_0\| \|F'(y_0)^{-1} F'(x^*)\| \right. \\ &\quad \left. + \|I - \frac{3}{2} J_0\| \|F'(x_0)^{-1} F'(x^*)\| \|F'(x^*)^{-1} F(z_0)\| \right] \\ &\leq \left\{ \frac{3}{4} \left(1 + \frac{1}{1 - g_3(\|x_0 - x^*\|)}\right) \frac{M}{1 - L_0 g_2(\|x_0 - x^*\|) \|x_0 - x^*\|} \right. \\ &\quad \left. + \left(1 + \frac{3}{4} \left(1 + \frac{1}{1 - g_3(\|x_0 - x^*\|)}\right)\right) \frac{M}{1 - L_0 g_2(\|x_0 - x^*\|) \|x_0 - x^*\|} \right\} \|z_0 - x^*\|, \end{aligned}$$

so, from the preceding estimate,

$$\begin{aligned} \|x_1 - x^*\| &\leq \|z_0 - x^*\| + \|A_0 F(z_0)\| \\ &\leq g_4(\|x_0 - x^*\|) \|x_0 - x^*\| + \|A_0 F(z_0)\| \\ &\leq \left\{ 1 + \frac{3}{4} \left(1 + \frac{1}{1 - g_3(\|x_0 - x^*\|)}\right) \frac{M}{1 - L_0 g_2(\|x_0 - x^*\|) \|x_0 - x^*\|} \right. \\ &\quad \left. + \left(1 + \frac{3}{4} \left(1 + \frac{1}{1 - g_3(\|x_0 - x^*\|)}\right)\right) \frac{M}{1 - L_0 \|x_0 - x^*\|} \right\} g_4(\|x_0 - x^*\|) \|x_0 - x^*\| \\ &= g_5(\|x_0 - x^*\|) \|x_0 - x^*\| \\ &< \|x_0 - x^*\| < r^*, \end{aligned}$$

(by (2.7)) which shows (2.17) for  $n = 0$ . By simply replacing  $u_0, y_0, z_0, x_1$  by  $u_k, y_k, z_k, x_{k+1}$  in the preceding estimates we arrive at estimates (2.13)-(2.17). Finally, from the estimates  $\|x_{k+1} - x^*\| < \|x_k - x^*\|$  we obtain  $\lim_{k \rightarrow \infty} x_k = x^*$ .  $\square$

**REMARK 2.2.** 1. In view of (2.10) and the estimate

$$\begin{aligned} \|F'(x^*)^{-1} F'(x)\| &= \|F'(x^*)^{-1} (F'(x) - F'(x^*)) + I\| \\ &\leq 1 + \|F'(x^*)^{-1} (F'(x) - F'(x^*))\| \leq 1 + L_0 \|x - x^*\| \end{aligned}$$

condition (2.12) can be dropped and  $M$  can be replaced by

$$M(t) = 1 + L_0 t.$$

Moreover, condition (2.11) can be replaced by the popular but stronger conditions

$$\|F'(x^*)^{-1} (F'(x) - F'(y))\| \leq L \|x - y\| \text{ for each } x, y \in D \quad (2.21)$$

or

$$\|F'(x^*)^{-1} (F'(x^* + t(x - x^*)) - F'(x))\| \leq L(1-t) \|x - x^*\| \text{ for each } x, y \in D \text{ and } t \in [0, 1].$$

2. The results obtained here can be used for operators  $F$  satisfying autonomous differential equations [3] of the form

$$F'(x) = P(F(x))$$

where  $P$  is a continuous operator. Then, since  $F'(x^*) = P(F(x^*)) = P(0)$ , we can apply the results without actually knowing  $x^*$ . For example, let  $F(x) = e^x - 1$ . Then, we can choose:  $P(x) = x + 1$ .

3. The local results obtained here can be used for projection methods such as the Arnoldi's method, the generalized minimum residual method (GMRES), the generalized conjugate method (GCR) for combined Newton/finite projection methods and in connection to the mesh independence principle can be used to develop the cheapest and most efficient mesh refinement strategies [3, 4].
4. The radius  $r_A$  given in (2.3) was shown by us to be the convergence radius of Newton's method [3, 4]

$$x_{n+1} = x_n - F'(x_n)^{-1}F(x_n) \text{ for each } n = 0, 1, 2, \dots \quad (2.22)$$

under the conditions (2.10) and (2.21). It follows from (2.3) and (2.8) that the convergence radius  $r^*$  of the three-step method (1.2) cannot be larger than the convergence radius  $r_A$  of the second order Newton's method (2.22). As already noted in [3, 4]  $r_A$  is at least as large as the convergence ball given by Rheinboldt [3, 4]

$$r_R = \frac{2}{3L}.$$

In particular, for  $L_0 < L$  we have that

$$r_R < r_A$$

and

$$\frac{r_R}{r_A} \rightarrow \frac{1}{3} \text{ as } \frac{L_0}{L} \rightarrow 0.$$

That is our convergence ball  $r_A$  is at most three times larger than Rheinboldt's. The same value for  $r_R$  was given by Traub [3, 4].

5. It is worth noticing that method (1.2) is not changing when we use the conditions of Theorem 2.1 instead of the stronger (C) conditions used in [26]. Moreover, we can compute the computational order of convergence (COC) defined by

$$\xi = \ln \left( \frac{\|x_{n+1} - x^*\|}{\|x_n - x^*\|} \right) / \ln \left( \frac{\|x_n - x^*\|}{\|x_{n-1} - x^*\|} \right)$$

or the approximate computational order of convergence

$$\xi_1 = \ln \left( \frac{\|x_{n+1} - x_n\|}{\|x_n - x_{n-1}\|} \right) / \ln \left( \frac{\|x_n - x_{n-1}\|}{\|x_{n-1} - x_{n-2}\|} \right).$$

This way we obtain in practice the order of convergence in a way that avoids the bounds given in [26] involving estimates up to the third Fréchet derivative of operator  $F$ .

### 3. NUMERICAL EXAMPLES

We present numerical examples in this section.

**EXAMPLE 3.1.** Let  $X = Y = \mathbb{R}^2$ ,  $D = \bar{U}(0, 1)$ ,  $x^* = (0, 0)^T$  and define function  $F$  on  $D$  by

$$F(x) = (\sin x, \frac{1}{3}(e^x + 2x - 1))^T.$$

Then, using (2.10)-(2.12), we get  $L_0 = L = 1$ ,  $M = \frac{1}{3}(e + 2)$ . Then, by (2.8) we obtain

$$r^* = 0.0270 < r_R = r_A = 0.6667$$



**EXAMPLE 3.2.** Let  $X = Y = \mathbb{R}^3$ ,  $D = \bar{U}(0, 1)$ . Define  $F$  on  $D$  for  $v = (x, y, z)^T$  by

$$F(v) = (e^x - 1, \frac{e-1}{2}y^2 + y, z)^T.$$

Then, the Fréchet-derivative is given by

$$F'(v) = \begin{bmatrix} e^x & 0 & 0 \\ 0 & (e-1)y + 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Notice that  $x^* = (0, 0, 0)^T$ ,  $F'(x^*) = F'(x^*)^{-1} = \text{diag}\{1, 1, 1\}$ ,  $L_0 = e - 1 < L = e$ ,  $M = e$ . Then, by (2.8) we obtain

$$r^* = 0.0045 < r_R = 0.2453 < r_A = 0.3249.$$

#### REFERENCES

- [1] F. Ahmad, S. Hussain, N.A. Mir and A. Rafiq, New sixth order Jarratt method for solving nonlinear equations, *Int. J. Appl. Math. Mech.*, **5**(5), (2009), 27-35.
- [2] S. Amat, M.A. Hernández and N. Romero, A modified Chebyshev's iterative method with at least sixth order of convergence, *Appl. Math. Comput.*, **206**(1), (2008), 164-174.
- [3] I.K. Argyros, *Convergence and Application of Newton-type Iterations*, Springer, 2008.
- [4] I. K. Argyros and Said Hilout, A convergence analysis for directional two-step Newton methods, *Numer. Algor.*, **55**, (2010), 503-528.
- [5] D.D. Bruns and J.E. Bailey, Nonlinear feedback control for operating a nonisothermal CSTR near an unstable steady state, *Chem. Eng. Sci.*, **32**, (1977), 257-264.
- [6] V. Candela and A. Marquina, Recurrence relations for rational cubic methods I: The Halley method, *Computing*, **44**, (1990), 169-184.
- [7] V. Candela and A. Marquina, Recurrence relations for rational cubic methods II: The Chebyshev method, *Computing*, **45**(4), (1990), 355-367.
- [8] C. Chun, Some improvements of Jarratt's method with sixth-order convergence, *Appl. Math. Comput.*, **190**(2), (1990), 1432-1437.
- [9] J. A. Ezquerro and M.A. Hernández, Recurrence relations for Chebyshev-type methods, *Appl. Math. Optim.*, **41**(2), (2000), 227-236.
- [10] J. A. Ezquerro and M.A. Hernández, New iterations of R-order four with reduced computational cost. *BIT Numer. Math.*, **49**, (2009), 325- 342.
- [11] J. A. Ezquerro and M.A. Hernández, On the R-order of the Halley method, *J. Math. Anal. Appl.*, **303**, (2005), 591-601.
- [12] J.M. Gutiérrez and M.A. Hernández, Recurrence relations for the super-Halley method, *Computers Math. Applic.*, **36**(7), (1998), 1-8.
- [13] M. Ganesh and M.C. Joshi, Numerical solvability of Hammerstein integral equations of mixed type, *IMA J. Numer. Anal.*, **11**, ( 1991), 21-31.
- [14] M.A. Hernández, Chebyshev's approximation algorithms and applications, *Computers Math. Applic.*, **41**(3-4), (2001), 433-455.
- [15] M.A. Hernández and M.A. Salanova, Sufficient conditions for semilocal convergence of a fourth order multipoint iterative method for solving equations in Banach spaces. *Southwest J. Pure Appl. Math*(1), (1999), 29-40.
- [16] P. Jarratt, Some fourth order multipoint methods for solving equations, *Math. Comput.*, **20**(95), (1966), 434-437.
- [17] J. Kou and Y. Li, An improvement of the Jarratt method, *Appl. Math. Comput.*, **189**,(2007), 1816-1821.
- [18] S. K. Parhi and D.K. Gupta, Semilocal convergence of a stirling-like method in Banach spaces, *Int. J. Comput. Methods*, **7**(02), (2010), 215-228.
- [19] S. K. Parhi and D.K. Gupta, Recurrence relations for a Newton-like method in Banach spaces, *J. Comput. Appl. Math.*, **206**(2), (2007), 873-887.
- [20] L. B. Rall, *Computational solution of nonlinear operator equations*, Robert E. Krieger, New York,(1979).
- [21] H. Ren, Q. Wu and W. Bi, New variants of Jarratt method with sixth-order convergence, *Numer. Algorithms*, **52**(4), (2009), 585-603.

- [22] X. Wang, J. Kou and Y. Li, Modified Jarratt method with sixth order convergence, Appl. Math. Lett., **22**, (2009), 1798-1802.
- [23] X. Ye and C. Li, Convergence of the family of the deformed Euler-Halley iterations under the Hölder condition of the second derivative, J. Comput. Appl. Math., **194**(2), (2006), 294-308.
- [24] X. Ye, C. Li and W. Shen, Convergence of the variants of the Chebyshev-Halley iteration family under the Hölder condition of the first derivative, J. Comput. Appl. Math., **203**(1), (2007), 279-288.
- [25] Y. Zhao and Q. Wu, Newton-Kantorovich theorem for a family of modified Halley's method under Hölder continuity condition in Banach spaces, Appl. Math. Comput. **202**(1), (2008), 243-251.
- [26] X. Wang, J. Kou and C. Gu, Semilocal convergence of a sixth-order Jarratt method in Banach spaces, Numer. Algorithms, **57**, (2011), 441-456.
- [27] J. Kou and X. Wang, Semilocal convergence of a modified multi-point Jarratt method in Banach spaces under general continuity conditions, Numer. Algorithms, **60**, (2012), 369-390.