



SOME INTEGRAL INEQUALITIES OF THE HADAMARD AND THE FEJÉR-HADAMARD TYPE VIA GENERALIZED FRACTIONAL INTEGRAL OPERATOR

GHULAM ABBAS¹ AND GHULAM FARID*²

¹ Department of Mathematics, Government College Bhalwal, Sargodha, Pakistan

² Department of Mathematics, COMSATS University Islamabad, Attock, Pakistan

ABSTRACT.

In this paper we give the Hadamard and the Fejér-Hadamard type integral inequalities for convex and relative convex functions by involving a generalization of the Riemann-Liouville fractional integral. Also some connections with known results have been obtained. **KEYWORDS:** Convex function; Hadamad inequality; Fejér-Hadamard inequality; Fractional integral operators.

AMS Subject Classification: Primary 26A51; Secondary 26A33, 33E12.

1. PRELIMINARIES

Convex functions are very useful for diverse fields of Mathematics, a rich literature has been built since their discovery [15].

Definition 1.1. Let I be an interval of real numbers. Then a function $f : I \rightarrow \mathbb{R}$ is said to be convex function if for all $x, y \in I$ and $0 \leq \lambda \leq 1$ the following inequality holds

$$f(x\lambda + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y).$$

Convex functions are naturally obey the following inequality which is well known as the Hadamard inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}$$

where $f : I \rightarrow \mathbb{R}$ is a convex function on I and $a, b \in I, a < b$.

Following definitions are given in [14].

* Corresponding author.

Email address : prof.abbas6581@gmail.com (Ghulam Abbas), faridphdms@hotmail.com, ghlmfarid@cuiatk.edu.pk (Ghulam Farid).

Definition 1.2. Let T_g be a set of real numbers. This set T_g is said to be relative convex with respect to an arbitrary function $g : \mathbb{R} \rightarrow \mathbb{R}$ if

$$(1-t)x + tg(y) \in T_g$$

where $x, y \in \mathbb{R}$ such that $x, g(y) \in T_g$, $0 \leq t \leq 1$.

Note that every convex set is relative convex, but the converse is not true. For example $T_g = [-1, \frac{-1}{2}] \cup [0, 1]$ and $g(x) = x^2$, for all $x \in \mathbb{R}$. This set is relative convex but not convex set. Another possibility may be occur that a relative convex set is convex set for example if $T_g = [-1, 1]$ and $g(x) = (|x|)^{\frac{1}{4}}$ for all $x \in \mathbb{R}$ (see[9]). If $g = I$ the identity function, then the definition of relative convex set recaptures the definition of classical convex set.

Definition 1.3. A function $f : T_g \rightarrow \mathbb{R}$ is said to be relative convex, if there exists an arbitrary function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$f((1-t)x + tg(y)) \leq (1-t)f(x) + tf(g(y)),$$

holds, where $x, y \in \mathbb{R}$ such that $x, g(y) \in T_g$, $0 \leq t \leq 1$.

Noor et al proved the following Hadamard type integral inequality in [14] for relative convex functions via Riemann-Liouville fractional integral operators.

Theorem 1.4. Let f be a positive relative convex function and integrable on $[a, g(b)]$. Then the following inequality holds

$$f\left(\frac{a+g(b)}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(g(b)-a)^\alpha} [I_{a^+}^\alpha f(g(b)) + I_{b^-}^\alpha f(a)] \leq \frac{f(a) + f(g(b))}{2}$$

$\alpha > 0$.

In the following we give some definitions and known facts about fractional integral operators [17].

Definition 1.5. Let $\omega \in \mathbb{R}$ and $\alpha, \beta, k, l, \gamma$ be positive real numbers. The generalized fractional integral operators $\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k}$ and $\epsilon_{\alpha, \beta, l, \omega, b^-}^{\gamma, \delta, k}$ for a real valued continuous function f are defined as follows

$$\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} f\right)(x) = \int_a^x (x-t)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega(x-t)^\alpha) f(t) dt, \quad (1.1)$$

and

$$\left(\epsilon_{\alpha, \beta, l, \omega, b^-}^{\gamma, \delta, k} f\right)(x) = \int_x^b (t-x)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega(t-x)^\alpha) f(t) dt,$$

where the function $E_{\alpha, \beta, l}^{\gamma, \delta, k}$ is the generalized Mittag-Leffler function defined as

$$E_{\alpha, \beta, l}^{\gamma, \delta, k}(t) = \sum_{n=0}^{\infty} \frac{(\gamma)_{kn} t^n}{\Gamma(\alpha n + \beta)(\delta)_{ln}}, \quad (1.2)$$

the Pochhammer symbol $(a)_n$ is defined by $(a)_n = a(a+1)(a+2)\dots(a+n-1)$, $(a)_0 = 1$.

For $\omega = 0$, (1.1) produces the definition of Riemann-Liouville fractional integral operators [17]

$$I_{a^+}^\beta f(x) = \frac{1}{\Gamma(\beta)} \int_a^x (x-t)^{\beta-1} f(t) dt, \quad x > a$$

and

$$I_{b^-}^\beta f(x) = \frac{1}{\Gamma(\beta)} \int_x^b (t-x)^{\beta-1} f(t) dt, \quad x < b.$$

In [17] properties of the generalized Mittag-Leffler function are discussed and it is given that $E_{\alpha,\beta,l}^{\gamma,\delta,k}(t)$ is absolutely convergent for $k < l + \alpha$. Let S be the sum of series of absolute terms of the Mittag-Leffler function $E_{\alpha,\beta,l}^{\gamma,\delta,k}(t)$, then we have $|E_{\alpha,\beta,l}^{\gamma,\delta,k}(t)| \leq S$. We use this property of Mittag-Leffler function in our results where we need.

In [10] the following Hadamard and the Fejér-Hadamard inequalities for convex functions via generalized fractional integral operator containing the Mittag-Leffler function have been proved.

Theorem 1.6. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a positive function with $0 \leq a < b$ and $f \in L_1[a, b]$. If f is convex on $[a, b]$, then the following inequality for generalized fractional integrals holds*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) (\epsilon_{\alpha,\beta,l,\omega',a^+}^{\gamma,\delta,k} 1)(b) &\leq \frac{(\epsilon_{\alpha,\beta,l,\omega',a^+}^{\gamma,\delta,k} f)(b) + (\epsilon_{\alpha,\beta,l,\omega',b^-}^{\gamma,\delta,k} f)(a)}{2} \\ &\leq \frac{f(a) + f(b)}{2} (\epsilon_{\alpha,\beta,l,\omega',b^-}^{\gamma,\delta,k} 1)(a), \end{aligned} \quad (1.3)$$

where $\omega' = \frac{w}{(b-a)^\alpha}$.

Theorem 1.7. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function with $0 \leq a < b$ and $f \in L_1[a, b]$. Also, let $g : [a, b] \rightarrow \mathbb{R}$ be a function which is non-negative, integrable and symmetric about $\frac{a+b}{2}$. Then the following inequality for generalized fractional integrals holds*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) (\epsilon_{\alpha,\beta,l,\omega',a^+}^{\gamma,\delta,k} g)(b) &\leq \frac{(\epsilon_{\alpha,\beta,l,\omega',a^+}^{\gamma,\delta,k} fg)(b) + (\epsilon_{\alpha,\beta,l,\omega',b^-}^{\gamma,\delta,k} fg)(a)}{2} \\ &\leq \frac{f(a) + f(b)}{2} (\epsilon_{\alpha,\beta,l,\omega',b^-}^{\gamma,\delta,k} g)(a), \end{aligned} \quad (1.4)$$

where $\omega' = \frac{w}{(b-a)^\alpha}$.

In [12, 14] the Hadamard and the Fejér-Hadamard type inequalities for convex and relative convex functions via Riemann-Liouville fractional integral operators have been proved. In this paper we give fractional integral inequalities of the Hadamard and the Fejér-Hadamard type for convex and relative convex functions by using the fractional integral operators involving the generalized Mittag-Leffler function. We also produce the results which are given in [12, 14] by setting particular values of parameters.

2. MAIN RESULTS

Following lemmas are useful to establish new results.

Lemma 2.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an integrable and symmetric function about $\frac{a+b}{2}$. Then the following equality holds*

$$\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} f\right)(b) = \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} f\right)(a) = \frac{\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} f\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} f\right)(a)}{2}. \quad (2.1)$$

Proof. As f is symmetric about $\frac{a+b}{2}$, therefore $f(a+b-t) = f(t)$. By definition we have

$$\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} f\right)(b) = \int_a^b (b-t)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-t)^\alpha) f(t) dt, \quad (2.2)$$

replacing t by $a + b - t$ in equation (2.2) we have

$$\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} f\right)(b) = \int_a^b (t-a)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(t-a)^\alpha) f(t) dt.$$

This implies

$$\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} f\right)(b) = \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} f\right)(a). \quad (2.3)$$

Therefore we get (2.1). \square

Lemma 2.2. *Let $f : [a,b] \rightarrow \mathbb{R}$ be a differentiable function on (a,b) and $f' \in L[a,b]$. If $g : [a,b] \rightarrow \mathbb{R}$ is integrable and symmetric about $\frac{a+b}{2}$, then we have the following equality*

$$\begin{aligned} & \left(\frac{f(a) + f(b)}{2}\right) \left[\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} g\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} g\right)(a)\right] \\ & - \left[\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} gf\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} gf\right)(a)\right] \\ & = \int_a^b \left[\int_a^t (b-s)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-s)^\alpha) g(s) ds \right. \\ & \quad \left. - \int_t^b (s-a)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(s-a)^\alpha) g(s) ds \right] f'(t) dt. \end{aligned}$$

Proof. To prove this lemma we take terms of the right hand side, on integrating by parts and after simplification we have

$$\begin{aligned} & \int_a^b \left[\int_a^t (b-s)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-s)^\alpha) g(s) ds \right] f'(t) dt \\ & = f(b) \int_a^b (b-s)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-s)^\alpha) g(s) ds - \int_a^b (b-t)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-t)^\alpha) gf(t) dt \\ & = f(b) \left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} g\right)(b) - \left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} gf\right)(b). \end{aligned}$$

By using Lemma 2.1 we have

$$\begin{aligned} & \int_a^b \left[\int_a^t (b-s)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(b-s)^\alpha) g(s) ds \right] f'(t) dt \\ & = \frac{f(b)}{2} \left[\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} g\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} g\right)(a)\right] - \left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} gf\right)(b). \end{aligned} \quad (2.4)$$

Similarly

$$\begin{aligned} & - \int_t^b \left[(s-a)^{\beta-1} E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega(s-a)^\alpha) g(s) ds \right] f'(t) dt \\ & = \frac{f(a)}{2} \left[\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} g\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} g\right)(a)\right] - \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} gf\right)(a). \end{aligned} \quad (2.5)$$

Adding (2.4) and (2.5) we get the left hand side. \square

In the following we give our first integral inequality of the Hadamard type.

Theorem 2.3. *Let $f : I \rightarrow \mathbb{R}$ be a differentiable mapping in the interior of I with $f' \in L[a,b]$, $a < b$. If $|f'|$ is convex on $[a,b]$ and $g : I \rightarrow \mathbb{R}$ is continuous and symmetric function about $\frac{a+b}{2}$, then we have the following inequality*

$$\left| \left(\frac{f(a) + f(b)}{2}\right) \left[\left(\epsilon_{\alpha,\beta,l,\omega,a^+}^{\gamma,\delta,k} g\right)(b) + \left(\epsilon_{\alpha,\beta,l,\omega,b^-}^{\gamma,\delta,k} g\right)(a)\right] \right|$$

$$\begin{aligned} & - \left[\left(\epsilon_{\alpha, \beta, l, \omega, a+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b-}^{\gamma, \delta, k} g f \right) (a) \right] \\ & \leq \frac{\|g\|_{\infty} S(b-a)^{\beta+1}}{\beta(\beta+1)} \left(1 - \frac{1}{2^{\beta}} \right) [|f'(a) + f'(b)|], \end{aligned}$$

for $k < l + \alpha$ and $\|g\|_{\infty} = \sup_{t \in [a, b]} |g(t)|$.

Proof. By using Lemma 2.2 we have

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left[\left(\epsilon_{\alpha, \beta, l, \omega, a+}^{\gamma, \delta, k} g \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b-}^{\gamma, \delta, k} g \right) (a) \right] \right. \\ & - \left. \left[\left(\epsilon_{\alpha, \beta, l, \omega, a+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b-}^{\gamma, \delta, k} g f \right) (a) \right] \right| \\ & \leq \int_a^b \left| \left[\int_a^t (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s) ds \right. \right. \\ & \left. \left. - \int_t^b (s-a)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(s-a)^{\alpha}) g(s) ds \right] |f'(t)| dt. \right. \end{aligned} \quad (2.6)$$

Using the convexity of $|f'|$ we have

$$|f'(t)| \leq \frac{b-t}{b-a} |f'(a)| + \frac{t-a}{b-a} |f'(b)|; \quad t \in [a, b]. \quad (2.7)$$

By using symmetry of function g we have

$$\begin{aligned} & \int_t^b (s-a)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(s-a)^{\alpha}) g(s) ds \\ & = \int_a^{a+b-t} (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(a+b-s) ds \\ & = \int_a^{a+b-t} (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s) ds. \end{aligned}$$

This implies

$$\begin{aligned} & \left| \int_a^t (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s) ds - \int_t^b (s-a)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(s-a)^{\alpha}) g(s) ds \right| \\ & = \left| \int_t^{a+b-t} (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s) ds \right| \\ & \leq \begin{cases} \int_t^{a+b-t} |(b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s)| ds, & t \in [a, \frac{a+b}{2}] \\ \int_{a+b-t}^t |(b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s)| ds, & t \in [\frac{a+b}{2}, b]. \end{cases} \end{aligned} \quad (2.8)$$

By (2.6), (2.7), (2.8) and absolute convergence of Mittag-Leffler function, we have

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) \left[\left(\epsilon_{\alpha, \beta, l, \omega, a+}^{\gamma, \delta, k} g \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b-}^{\gamma, \delta, k} g \right) (a) \right] \right. \\ & - \left. \left[\left(\epsilon_{\alpha, \beta, l, \omega, a+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b-}^{\gamma, \delta, k} g f \right) (a) \right] \right| \\ & \leq \int_a^{\frac{a+b}{2}} \left(\int_a^{a+b-t} |(b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s)| ds \right) \left(\frac{b-t}{b-a} |f'(a)| + \frac{t-a}{b-a} |f'(b)| \right) dt \\ & + \int_{\frac{a+b}{2}}^b \left(\int_{a+b-t}^t |(b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^{\alpha}) g(s)| ds \right) \left(\frac{b-t}{b-a} |f'(a)| + \frac{t-a}{b-a} |f'(b)| \right) dt. \end{aligned} \quad (2.9)$$

$$\begin{aligned} &\leq \frac{\|g\|_\infty S}{\beta(b-a)} \left[\int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta(b-t)|f'(a)|)dt + \int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta(t-a)|f'(b)|)dt \right. \\ &\quad \left. + \int_{\frac{a+b}{2}}^b ((t-a)^\beta - (b-t)^\beta(b-t)|f'(a)|)dt + \int_{\frac{a+b}{2}}^b ((t-a)^\beta - (b-t)^\beta(t-a)|f'(b)|)dt \right]. \end{aligned}$$

Since we have

$$\int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta)(b-t)dt = \frac{(b-a)^{\beta+2}}{\beta+1} \left(\frac{\beta+1}{\beta+2} - \frac{1}{2^{\beta+1}} \right)$$

and

$$\int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta)(t-a)dt = \frac{(b-a)^{\beta+2}}{\beta+1} \left(\frac{1}{\beta+2} - \frac{1}{2^{\beta+1}} \right).$$

Using the above calculations in (2.9) we have

$$\begin{aligned} &\left| \left(\frac{f(a)+f(b)}{2} \right) \left[\left(\epsilon_{\alpha,\beta,l,\omega,a+g}^{\gamma,\delta,k} \right) (b) + \left(\epsilon_{\alpha,\beta,l,\omega,b-g}^{\gamma,\delta,k} \right) (a) \right] \right. \\ &\quad \left. - \left[\left(\epsilon_{\alpha,\beta,l,\omega,a+gf}^{\gamma,\delta,k} \right) (b) + \left(\epsilon_{\alpha,\beta,l,\omega,b-gf}^{\gamma,\delta,k} \right) (a) \right] \right| \\ &\leq \frac{\|g\|_\infty S}{\beta(b-a)} \frac{(b-a)^{\beta+2}}{\beta+1} \left[\left(\frac{\beta+1}{\beta+2} - \frac{1}{2^{\beta+1}} \right) + \left(\frac{1}{\beta+2} - \frac{1}{2^{\beta+1}} \right) \right] [|f'(a)| + |f'(b)|] \\ &= \frac{\|g\|_\infty S}{\beta(\beta+1)} (b-a)^{\beta+1} \left(1 - \frac{1}{2^\beta} \right) [|f'(a)| + |f'(b)|]. \end{aligned}$$

□

A special case is stated in the following, which is inequality of the Hadamard type for Riemann-Liouville fractional integrals.

Corollary 2.4. *Setting $\omega = 0$ in Theorem 2.3 we have the following inequality for Riemann-Liouville fractional integral operators*

$$\begin{aligned} &\left| \left(\frac{f(a)+f(b)}{2} \right) \left[I_{a+}^\beta g(b) + I_{b-}^\beta g(a) \right] - \left[I_{a+}^\beta fg(b) + I_{b-}^\beta fg(a) \right] \right| \quad (2.10) \\ &\leq \frac{\|g\|_\infty (b-a)^{\beta+1}}{\Gamma(\beta+2)} \left(1 - \frac{1}{2^\beta} \right) [|f'(a)| + |f'(b)|]. \end{aligned}$$

Remark 2.5. The above inequality (2.10) is proved in [12].

Theorem 2.6. *Let $f : I \rightarrow \mathbb{R}$ be a differentiable function in the interior of I , also let $f' \in L[a, b]$, $a < b$. If $|f'|^q$, $q > 0$ is convex on $[a, b]$ and $g : I \rightarrow \mathbb{R}$ is continuous and symmetric function about $\frac{a+b}{2}$, then we have the following inequality*

$$\begin{aligned} &\left| \left(\frac{f(a)+f(b)}{2} \right) \left[\left(\epsilon_{\alpha,\beta,l,\omega,a+g}^{\gamma,\delta,k} \right) (b) + \left(\epsilon_{\alpha,\beta,l,\omega,b-g}^{\gamma,\delta,k} \right) (a) \right] \right. \\ &\quad \left. - \left[\left(\epsilon_{\alpha,\beta,l,\omega,a+gf}^{\gamma,\delta,k} \right) (b) + \left(\epsilon_{\alpha,\beta,l,\omega,b-gf}^{\gamma,\delta,k} \right) (a) \right] \right| \quad (2.11) \\ &\leq \frac{2\|g\|_\infty S (b-a)^{\beta+\frac{1}{p}}}{\beta(\beta+1)} \left(1 - \frac{1}{2^\beta} \right) (|f'(a)|^q + |f'(b)|^q)^{\frac{1}{q}}, \end{aligned}$$

for $k < l + \alpha$ and $\|g\|_\infty = \sup_{t \in [a,b]} |g(t)|$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 2.2, Hölder inequality, inequality (2.8) one can has

$$\left| \left(\frac{f(a)+f(b)}{2} \right) \left[\left(\epsilon_{\alpha,\beta,l,\omega,a+g}^{\gamma,\delta,k} \right) (b) + \left(\epsilon_{\alpha,\beta,l,\omega,b-g}^{\gamma,\delta,k} \right) (a) \right] \right| \quad (2.12)$$

$$\begin{aligned}
& - \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g f \right) (a) \right] \\
& \leq \left[\int_a^b \left| \int_t^{a+b-t} (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^\alpha) g(s) ds \right| dt \right]^{1-\frac{1}{q}} \\
& \quad \left[\int_a^b \left| \int_t^{a+b-t} (b-s)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} (\omega(b-s)^\alpha) g(s) ds \right| |f'(t)|^q dt \right]^{\frac{1}{q}}.
\end{aligned}$$

Using absolute convergence of Mittag-Leffler function and $\|g\|_\infty = \sup_{t \in [a, b]} |g(t)|$ we

have

$$\begin{aligned}
& \left| \left(\frac{f(a) + f(b)}{2} \right) \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g \right) (a) \right] \right. \\
& \quad \left. - \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g f \right) (a) \right] \right| \\
& \leq \|g\|_\infty^{1-\frac{1}{q}} S^{1-\frac{1}{q}} \left[\int_a^{\frac{a+b}{2}} \left(\int_t^{a+b-t} (b-s)^{\beta-1} ds \right) dt + \int_{\frac{a+b}{2}}^b \left(\int_{a+b-t}^t (b-s)^{\beta-1} ds \right) dt \right]^{1-\frac{1}{q}} \\
& \times \|g\|_\infty^{\frac{1}{q}} S^{\frac{1}{q}} \left[\int_a^{\frac{a+b}{2}} \left(\int_t^{a+b-t} (b-s)^{\beta-1} ds \right) |f'(t)|^q dt \right. \\
& \quad \left. + \int_{\frac{a+b}{2}}^b \left(\int_{a+b-t}^t (b-s)^{\beta-1} ds \right) |f'(t)|^q dt \right]^{\frac{1}{q}}.
\end{aligned}$$

By some calculation we have

$$\begin{aligned}
& \left| \left(\frac{f(a) + f(b)}{2} \right) \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g \right) (a) \right] \right. \\
& \quad \left. - \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g f \right) (a) \right] \right| \\
& \leq \|g\|_\infty S \left[\frac{(b-a)^{\beta+1}}{\beta+1} \left(1 - \frac{1}{2^\beta}\right) + \frac{(b-a)^{\beta+1}}{\beta+1} \left(1 - \frac{1}{2^\beta}\right) \right]^{1-\frac{1}{q}} \\
& \times \left[\int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta) |f'(t)|^q dt + \int_{\frac{a+b}{2}}^b ((b-t)^\beta - (t-a)^\beta) |f'(t)|^q dt \right]^{\frac{1}{q}}.
\end{aligned}$$

Since $|f'|^q$ is convex on $[a, b]$, therefore we have

$$|f'(t)|^q \leq \frac{b-t}{b-a} |f'(a)|^q + \frac{t-a}{b-a} |f'(b)|^q. \quad (2.13)$$

Hence

$$\begin{aligned}
& \left| \left(\frac{f(a) + f(b)}{2} \right) \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g \right) (a) \right] \right. \\
& \quad \left. - \left[\left(\epsilon_{\alpha, \beta, l, \omega, a^+}^{\gamma, \delta, k} g f \right) (b) + \left(\epsilon_{\alpha, \beta, l, \omega, b_-}^{\gamma, \delta, k} g f \right) (a) \right] \right| \\
& \leq \|g\|_\infty S \left[2 \frac{(b-a)^{\beta+1}}{\beta+1} \left(1 - \frac{1}{2^\beta}\right) \right]^{1-\frac{1}{q}} \\
& \times \left[\int_a^{\frac{a+b}{2}} ((b-t)^\beta - (t-a)^\beta) \left(\frac{b-t}{b-a} |f'(a)|^q + \frac{t-a}{b-a} |f'(b)|^q \right) dt \right.
\end{aligned}$$

$$+ \int_{\frac{a+b}{2}}^b ((b-t)^\beta - (t-a)^\beta) \left(\frac{b-t}{b-a} |f'(a)|^q + \frac{t-a}{b-a} |f'(b)|^q \right) dt \Bigg]^{\frac{1}{q}}.$$

From which one can have (2.11). \square

Corollary 2.7. *Setting $\omega = 0$ in Theorem 2.6 we have the following result for Riemann-Liouville fractional integral operators*

$$\begin{aligned} & \left| \left(\frac{f(a) + f(b)}{2} \right) [I_{a+}^\beta g(b) + I_{b-}^\beta g(a)] - [I_{a+}^\beta f g(b) + I_{b-}^\beta f g(a)] \right| \\ & \leq \frac{2 \|g\|_\infty (b-a)^{\beta+1-\frac{1}{q}}}{\Gamma(\beta+2)} \left(1 - \frac{1}{2^\beta} \right) (|f'(a)|^q + |f'(b)|^q)^{\frac{1}{q}}, \end{aligned}$$

$\beta > 0$.

In the following we give the Hadamard inequality for relative convex functions via generalized fractional integral operators.

Theorem 2.8. *Let $f : [a, g(b)] \rightarrow \mathbb{R}$ be a positive relative convex function and $f \in L[a, g(b)]$. Then the following inequalities for generalized fractional integral operators hold*

$$\begin{aligned} f \left(\frac{a+g(b)}{2} \right) \left(\epsilon_{\alpha, \beta, l, \omega', a+1}^{\gamma, \delta, k} \right) (g(b)) & \leq \frac{1}{2} \left[\left(\epsilon_{\alpha, \beta, l, \omega', a+1}^{\gamma, \delta, k} f \right) (g(b)) + \left(\epsilon_{\alpha, \beta, l, \omega', g(b)-}^{\gamma, \delta, k} f \right) (a) \right] \\ & \leq \frac{f(a) + f(g(b))}{2} \left(\epsilon_{\alpha, \beta, l, \omega', g(b)-}^{\gamma, \delta, k} 1 \right) (a), \end{aligned}$$

where $\omega' = \frac{\omega}{(g(b)-a)^\alpha}$.

Proof. Since f is relative convex on $[a, g(b)]$, we have

$$\begin{aligned} f \left(\frac{a+g(b)}{2} \right) & = f \left[\left(\frac{1}{2} (ta + (1-t)g(b)) \right) + \left(1 - \frac{1}{2} \right) ((1-t)a + tg(b)) \right] \\ & \leq \frac{1}{2} f (ta + (1-t)g(b)) + \frac{1}{2} f ((1-t)a + tg(b)). \end{aligned}$$

Multiplying both sides by $2t^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega t^\alpha)$ and integrating over $[0, 1]$ we have

$$\begin{aligned} 2f \left(\frac{a+g(b)}{2} \right) \int_0^1 t^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega t^\alpha) dt & \leq \int_0^1 t^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega t^\alpha) f (ta + (1-t)g(b)) dt \\ & \quad + \int_0^1 t^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k}(\omega t^\alpha) f ((1-t)a + tg(b)) dt. \end{aligned} \quad (2.14)$$

Setting $ta + (1-t)g(b) = x$ that is $t = \frac{g(b)-x}{g(b)-a}$ and $(1-t)a + tg(b) = y$ that is $t = \frac{y-a}{g(b)-a}$ we have

$$\begin{aligned} & 2f \left(\frac{a+g(b)}{2} \right) \int_{g(b)}^a \left(\frac{g(b)-x}{g(b)-a} \right)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} \left(\omega \left(\frac{g(b)-x}{g(b)-a} \right)^\alpha \right) \left(\frac{-dx}{g(b)-a} \right) \\ & \leq \int_{g(b)}^a \left(\frac{g(b)-x}{g(b)-a} \right)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} \left(\omega \left(\frac{g(b)-x}{g(b)-a} \right)^\alpha \right) f(x) \left(\frac{-dx}{g(b)-a} \right) \\ & \quad + \int_a^{g(b)} \left(\frac{y-a}{g(b)-a} \right)^{\beta-1} E_{\alpha, \beta, l}^{\gamma, \delta, k} \left(\omega \left(\frac{y-a}{g(b)-a} \right)^\alpha \right) f(y) \left(\frac{dy}{g(b)-a} \right). \end{aligned} \quad (2.15)$$

After simplification we get

$$2f\left(\frac{a+g(b)}{2}\right)\left(\epsilon_{\alpha,\beta,l,\omega',a+}^{\gamma,\delta,k}\right)(g(b)) \leq \left[\left(\epsilon_{\alpha,\beta,l,\omega',a+}^{\gamma,\delta,k}\right)(g(b)) + \left(\epsilon_{\alpha,\beta,l,\omega',g(b)-}^{\gamma,\delta,k}\right)(a)\right]. \quad (2.16)$$

By using the relative convexity of f on $[a, g(b)]$ one can has

$$f(ta+(1-t)g(b)) + f((1-t)a+tg(b)) \leq tf(a) + (1-t)f(g(b)) + (1-t)f(a) + tf(g(b)).$$

Multiplying $t^{\beta-1}E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega t^\alpha)$ on both sides and integrating over $[0, 1]$ we have

$$\begin{aligned} & \int_0^1 t^{\beta-1}E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega t^\alpha)f(ta+(1-t)g(b))dt + \int_0^1 t^{\beta-1}E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega t^\alpha)f((1-t)a+tg(b))dt \\ & \leq \int_0^1 t^{\beta-1}E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega t^\alpha)(tf(a)+(1-t)f(g(b)))dt + \int_0^1 t^{\beta-1}E_{\alpha,\beta,l}^{\gamma,\delta,k}(\omega t^\alpha)((1-t)f(a)+tf(g(b)))dt. \end{aligned}$$

Setting $ta+(1-t)g(b) = x$ that is $t = \frac{g(b)-x}{g(b)-a}$ and $(1-t)a+tg(b) = y$ that is $t = \frac{y-a}{g(b)-a}$ and after simple calculation we have

$$\left[\left(\epsilon_{\alpha,\beta,l,\omega',a+}^{\gamma,\delta,k}\right)(g(b)) + \left(\epsilon_{\alpha,\beta,l,\omega',g(b)-}^{\gamma,\delta,k}\right)(a)\right] \leq [f(a)+f(g(b))]\left(\epsilon_{\alpha,\beta,l,\omega',g(b)-}^{\gamma,\delta,k}\right)(a). \quad (2.17)$$

Combinig (2.16) and (2.17) we get the result. \square

Remark 2.9. (i) If we put $\omega = 0$ and $k = 1$ in Theorem 2.8 we obtain Theorem 1.4.

(ii) If we put $\omega = 0$ and $\beta = \frac{\alpha}{k}$ in Theorem 2.8, then we get [11, Theorem 3].

In the upcoming theorem we give the generalization of previous result.

Theorem 2.10. *Let $f : [g(a), g(b)] \rightarrow \mathbb{R}$ be a positive relative convex function and $f \in L[g(a), g(b)]$. Then the following inequalities for generalized fractional integral operator holds*

$$\begin{aligned} & f\left(\frac{g(a)+g(b)}{2}\right)\left(\epsilon_{\alpha,\beta,l,\omega',g(a)+}^{\gamma,\delta,k}\right)(g(b)) \\ & \leq \frac{1}{2}\left[\left(\epsilon_{\alpha,\beta,l,\omega',g(a)+}^{\gamma,\delta,k}\right)(g(b)) + \left(\epsilon_{\alpha,\beta,l,\omega',g(b)-}^{\gamma,\delta,k}\right)(a)\right] \\ & \leq \frac{f(g(a))+f(g(b))}{2}\left(\epsilon_{\alpha,\beta,l,\omega',g(b)-}^{\gamma,\delta,k}\right)(g(a)), \end{aligned}$$

where $\omega' = \frac{\omega}{(g(b)-g(a))^\alpha}$.

Proof. Proof of this theorem is on the same lines of the proof of Theorem 2.8. \square

Corollary 2.11. *For $\omega = 0$ we obtain the following inequality for Riemann-Liouville integral operator from Theorem 2.10*

$$\begin{aligned} f\left(\frac{g(a)+g(b)}{2}\right) & \leq \frac{\Gamma(\beta+1)}{2(g(b)-g(a))^\beta}[I_{g(a)+}^\beta f(g(b)) + I_{g(b)-}^\beta f(g(a))] \\ & \leq \frac{f(g(a))+f(g(b))}{2}, \end{aligned}$$

with $\beta > 0$.

Remark 2.12. In Theorem 2.10 if we take $\omega = 0$, $\beta = \frac{\alpha}{k}$, then we get [11, Theorem 5].

3. ACKNOWLEDGMENTS

The research work of Ghulam Farid is supported by Higher Education Commission of Pakistan under NRP 2016, Project No. 5421.

REFERENCES

1. M. Adil Khan, Y. Khurshid, T. Ali, and N. Rehman, Inequalities for three times differentiable functions, *Punjab Univ. J. Math.*, 2016, **48**(2), 35-48.
2. M. Adil Khan, T. Ali, S. S. Dragomir, Hermite-Hadamard type inequalities for conformable fractional integrals, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math.*, (2017), DOI 10.1007/s13398-0170408-5.
3. M. Adil Khan, Y. Khurshid and T. Ali, Hermite-Hadamard inequality for fractional integrals Via η -convex functions, *Acta Math. Univ. Comenian.*, **86**(1) (2017), 153-164.
4. M. Adil Khan, Yu-Ming Chu, A. Kashuri, R. Liko, G. Ali, New Hermite-Hadamard inequalities for conformable fractional integrals, *Journal of Function spaces*, to appear.
5. M. Adil Khan, T. Ali, M. Z. Sarikaya, and Q. Din, New bounds for Hermite-Hadamard type inequalities with applications, *Electronic Journal of Mathematical Analysis and Applications*, to appear.
6. Y. M. Chu, M. Adil Khan, T. U. Khan, T. Ali, Generalizations of Hermite-Hadamard type inequalities for MT-convex functions, *J. Nonlinear Sci. Appl.*, **9** (2016), 4305-4316.
7. Y. M. Chu, M. Adil Khan, T. Ali, S. S. Dragomir, Inequalities for α -fractional differentiable functions, *J. Inequal. Appl.*, **2017** (2017), Article ID 93, 12 pages.
8. Y. M. Chu, M. Adil Khan, T. U. Khan, and J. Khan, Some new inequalities of Hermite-Hadamard type for s -convex functions with applications, *Open Math.*, **15** (2017) 1414-1430.
9. D. I. Duca, L. Lupa, Saddle points for vector valued functions: existence, necessary and sufficient theorems, *J. Glob. Optimization* **53** (2012), 431-440.
10. G. Farid, Hadamard and Fejér-Hadamard inequalities for generalized fractional integrals involving special functions, *Konuralp J. Math.* **4**(1) (2016), 108-113.
11. G. Farid, A. U. Rehman and M. Zahra, On Hadamard inequalities for relative convex functions via fractional integrals, *Nonlinear Anal. Forum* **21**(1) (2016) 77-86.
12. I. Iscan, Hermite Hadamard Fejér type inequalities for convex functions via fractional integrals, *Stud. Univ. Babeş-Bolyai Math.* **60**(3) (2015), 355-366.
13. M. A. Noor, Differential non-convex functions and general variational inequalities, *Appl. Math. Comp.*, **199**(2) (2008), 623-630.
14. M. A. Noor, K. I. Noor and M. U. Awan, *Generalized convexity and integral inequalities*, *Appl. Math. Inf. Sci.*, **9** (1) (2015), 233-243.
15. J. Pečarić, F. Proschan and Y. L. Tong, *Convex functions, partial orderings and statistical applications*, Academic Press, New York, 1992.
16. T. R. Prabhakar, A singular integral equation with a generalized Mittag-Leffler function in the kernel, *Yokohama Math. J.* **19** (1971), 7-15.
17. L. T. O. Salim and A. W. Faraj, A Generalization of Mittag-Leffler function and integral operator associated with integral calculus, *J. Frac. Calc. Appl.* **3**(5) (2012), 1-13.
18. E. Set, S. S. Karatas and M. Adil Khan, Hermite-Hadamard type inequalities obtained via fractional integral for differentiable m -convex and (α, m) -convex function, *International Journal of Analysis*, **2016**, Article ID 4765691, 8 pages.
19. H. M. Srivastava and Z. Tomovski, Fractional calculus with an integral operator containing generalized Mittag-Leffler function in the kernel, *Appl. Math. Comput.* **211**(1) (2009), 198-210.